

ON THE GEOMETRY OF BIHARMONIC SUBMANIFOLDS IN SASAKIAN SPACE FORMS

D. FETCU AND C. ONICIUC

ABSTRACT. We classify all proper-biharmonic Legendre curves in a Sasakian space form and point out some of their geometric properties. Then we provide a method for constructing anti-invariant proper-biharmonic submanifolds in Sasakian space forms. Finally, using the Boothby-Wang fibration, we determine all proper-biharmonic Hopf cylinders over homogeneous real hypersurfaces in complex projective spaces.

1. INTRODUCTION

As defined by Eells and Sampson in [14], *harmonic maps* $f : (M, g) \rightarrow (N, h)$ are the critical points of the *energy functional*

$$E(f) = \frac{1}{2} \int_M \|df\|^2 v_g$$

and they are solutions of the associated Euler-Lagrange equation

$$\tau(f) = \text{trace}_g \nabla df = 0,$$

where $\tau(f)$ is called the *tension field* of f . When f is an isometric immersion with mean curvature vector field H , then $\tau(f) = mH$ and f is harmonic if and only if it is minimal.

The *bienergy functional* (proposed also by Eells and Sampson in 1964, [14]) is defined by

$$E_2(f) = \frac{1}{2} \int_M \|\tau(f)\|^2 v_g.$$

The critical points of E_2 are called *biharmonic maps* and they are solutions of the Euler-Lagrange equation (derived by Jiang in 1986, [20]):

$$\tau_2(f) = -\Delta^f \tau(f) - \text{trace}_g R^N(df, \tau(f))df = 0,$$

where Δ^f is the Laplacian on sections of $f^{-1}TN$ and $R^N(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]}$ is the curvature operator on N ; $\tau_2(f)$ is called the *bitension field* of f . Since all harmonic maps are biharmonic, we are interested in studying those which are biharmonic but non-harmonic, called *proper-biharmonic maps*.

Now, if $f : M \rightarrow N_c$ is an isometric immersion into a space form of constant sectional curvature c , then

$$\tau(f) = mH \quad \text{and} \quad \tau_2(f) = -m\Delta^f H + cm^2H.$$

Thus f is biharmonic if and only if

$$\Delta^f H = mcH.$$

Contribution to the Proceedings of the 10-th International Conference on Geometry, Integrability and Quantization, Varna 2008, Bulgaria.

In a different way, Chen defined the biharmonic submanifolds in an Euclidean space as those with harmonic mean curvature vector field ([10]). Replacing $c = 0$ in the above equation we just reobtain Chen's definition. Moreover, let $f : M \rightarrow \mathbb{R}^n$ be an isometric immersion. Set $f = (f^1, \dots, f^n)$ and $H = (H^1, \dots, H^n)$. Then $\Delta^f H = (\Delta H^1, \dots, \Delta H^n)$, where Δ is the Beltrami-Laplace operator on M , and f is biharmonic if and only if

$$\Delta^f H = \Delta\left(\frac{-\Delta f}{m}\right) = -\frac{1}{m}\Delta^2 f = 0.$$

There are several classification results for the proper-biharmonic submanifolds in Euclidean spheres and non-existence results for such submanifolds in space forms N_c , $c \leq 0$ ([4], [5], [7], [8], [9], [10], [13]), while in spaces of non-constant sectional curvature only few results were obtained ([1], [12], [18], [19], [25], [29]).

We recall that the proper-biharmonic curves of the unit Euclidean 2-dimensional sphere $\mathbb{S}^2$ are the circles of radius $\frac{1}{\sqrt{2}}$, and the proper-biharmonic curves of $\mathbb{S}^3$ are the geodesics of the minimal Clifford torus $\mathbb{S}^1(\frac{1}{\sqrt{2}}) \times \mathbb{S}^1(\frac{1}{\sqrt{2}})$ with the slope different from ± 1 . The proper-biharmonic curves of $\mathbb{S}^3$ are helices. Further, the proper-biharmonic curves of $\mathbb{S}^n$, $n > 3$, are those of $\mathbb{S}^3$ (up to a totally geodesic embedding). Concerning the hypersurfaces of $\mathbb{S}^n$, it was conjectured in [4] that the only proper-biharmonic hypersurfaces are the open parts of $\mathbb{S}^{n-1}(\frac{1}{\sqrt{2}})$ or $\mathbb{S}^{m_1}(\frac{1}{\sqrt{2}}) \times \mathbb{S}^{m_2}(\frac{1}{\sqrt{2}})$ with $m_1 + m_2 = n - 1$ and $m_1 \neq m_2$.

Since odd dimensional unit Euclidean spheres $\mathbb{S}^{2n+1}$ are Sasakian space forms with constant φ -sectional curvature 1, the next step is to study the biharmonic submanifolds of Sasakian space forms. In this paper we mainly gather the results obtained in [15], [16] and [17].

We note that the proper-biharmonic submanifolds in pseudo-Riemannian manifolds are also intensively-studied (for example, see [2], [3], [11]).

For a general account of biharmonic maps see [22] and *The Bibliography of Biharmonic Maps* [28].

Conventions. We work in the C^∞ category, that means manifolds, metrics, connections and maps are smooth. The Lie algebra of the vector fields on N is denoted by $C(TN)$.

2. SASAKIAN SPACE FORMS

In this section we briefly recall some basic facts from the theory of Sasakian manifolds. For more details see [6].

A *contact metric structure* on a manifold N^{2n+1} is given by (φ, ξ, η, g) , where φ is a tensor field of type $(1, 1)$ on N , ξ is a vector field on N , η is a 1-form on N and g is a Riemannian metric, such that

$$\begin{cases} \varphi^2 = -I + \eta \otimes \xi, & \eta(\xi) = 1, \\ g(\varphi X, \varphi Y) = g(X, Y) - \eta(X)\eta(Y), & g(X, \varphi Y) = d\eta(X, Y), \end{cases}$$

for any $X, Y \in C(TN)$.

A contact metric structure (φ, ξ, η, g) is *Sasakian* if it is *normal*, i.e.

$$N_\varphi + 2d\eta \otimes \xi = 0,$$

where

$$N_\varphi(X, Y) = [\varphi X, \varphi Y] - \varphi[\varphi X, Y] - \varphi[X, \varphi Y] + \varphi^2[X, Y], \quad \forall X, Y \in C(TN)$$

is the Nijenhuis tensor field of φ .

The *contact distribution* of a Sasakian manifold $(N, \varphi, \xi, \eta, g)$ is defined by $\{X \in TN : \eta(X) = 0\}$, and an integral curve of the contact distribution is called *Legendre curve*.

A submanifold M of N which is tangent to ξ is said to be *anti-invariant* if φ maps any vector tangent to M and normal to ξ to a vector normal to M .

Let $(N, \varphi, \xi, \eta, g)$ be a Sasakian manifold. The sectional curvature of a 2-plane generated by X and φX , where X is a unit vector orthogonal to ξ , is called φ -*sectional curvature* determined by X . A Sasakian manifold with constant φ -sectional curvature c is called a *Sasakian space form* and it is denoted by $N(c)$.

A contact metric manifold $(N, \varphi, \xi, \eta, g)$ is called *regular* if for any point $p \in N$ there exists a cubic neighborhood of p such that any integral curve of ξ passes through the neighborhood at most once, and *strictly regular* if all integral curves are homeomorphic to each other.

Let $(N, \varphi, \xi, \eta, g)$ be a regular contact metric manifold. Then the orbit space $\bar{N} = N/\xi$ has a natural manifold structure and, moreover, if N is compact then N is a principal circle bundle over $\bar{N}$ (the Boothby-Wang Theorem). In this case the fibration $\pi : N \rightarrow \bar{N}$ is called *the Boothby-Wang fibration*. The Hopf fibration $\pi : \mathbb{S}^{2n+1} \rightarrow \mathbb{C}P^n$ is a well-known example of a Boothby-Wang fibration.

Theorem 2.1 ([24]). *Let $(N, \varphi, \xi, \eta, g)$ be a strictly regular Sasakian manifold. Then on $\bar{N}$ can be given the structure of a Kähler manifold. Moreover, if $(N, \varphi, \xi, \eta, g)$ is a Sasakian space form $N(c)$, then $\bar{N}$ has constant sectional holomorphic curvature $c + 3$.*

Even if N is non-compact, we still call the fibration $\pi : N \rightarrow \bar{N}$ of a strictly regular Sasakian manifold, the Boothby-Wang fibration.

3. BIHARMONIC LEGENDRE CURVES IN SASAKIAN SPACE FORMS

Let (N^n, g) be a Riemannian manifold and $\gamma : I \rightarrow N$ a curve parametrized by arc length. Then γ is called a *Frenet curve of osculating order r* , $1 \leq r \leq n$, if there exists orthonormal vector fields $E_1, E_2, \dots, E_r$ along γ such that $E_1 = \gamma' = T$, $\nabla_T E_1 = \kappa_1 E_2$, $\nabla_T E_2 = -\kappa_1 E_1 + \kappa_2 E_3, \dots, \nabla_T E_r = -\kappa_{r-1} E_{r-1}$, where $\kappa_1, \dots, \kappa_{r-1}$ are positive functions on I .

A geodesic is a Frenet curve of osculating order 1; a *circle* is a Frenet curve of osculating order 2 with $\kappa_1 = \text{constant}$; a *helix of order r* , $r \geq 3$, is a Frenet curve of osculating order r with $\kappa_1, \dots, \kappa_{r-1}$ constants; a helix of order 3 is called, simply, helix.

In [16] we studied the biharmonicity of Legendre Frenet curves and we obtained the following results.

Let $(N^{2n+1}, \varphi, \xi, \eta, g)$ be a Sasakian space form with constant φ -sectional curvature c and $\gamma : I \rightarrow N$ a Legendre Frenet curve of osculating order r . Then γ is

biharmonic if and only if

$$\begin{aligned}
\tau_2(\gamma) &= \nabla_T^3 T - R(T, \nabla_T T)T \\
&= (-3\kappa_1\kappa_1')E_1 + \left(\kappa_1'' - \kappa_1^3 - \kappa_1\kappa_2^2 + \frac{(c+3)\kappa_1}{4}\right)E_2 \\
&\quad + (2\kappa_1'\kappa_2 + \kappa_1\kappa_2')E_3 + \kappa_1\kappa_2\kappa_3E_4 + \frac{3(c-1)\kappa_1}{4}g(E_2, \varphi T)\varphi T \\
&= 0.
\end{aligned}$$

The expression of the bitension field $\tau_2(\gamma)$ imposed a case-by-case analysis as follows.

Case I ($c = 1$)

Theorem 3.1 ([16]). *If $c = 1$ then γ is proper-biharmonic if and only if $n \geq 2$ and either γ is a circle with $\kappa_1 = 1$ or γ is a helix with $\kappa_1^2 + \kappa_2^2 = 1$.*

Case II ($c \neq 1$ and $E_2 \perp \varphi T$)

Theorem 3.2 ([16]). *Assume that $c \neq 1$ and $E_2 \perp \varphi T$. We have*

- 1) *if $c \leq -3$ then γ is biharmonic if and only if it is a geodesic;*
- 2) *if $c > -3$ then γ is proper-biharmonic if and only if either*
 - a) *$n \geq 2$ and γ is a circle with $\kappa_1^2 = \frac{c+3}{4}$, or*
 - b) *$n \geq 3$ and γ is a helix with $\kappa_1^2 + \kappa_2^2 = \frac{c+3}{4}$.*

Case III ($c \neq 1$ and $E_2 \parallel \varphi T$)

Theorem 3.3 ([16]). *If $c \neq 1$ and $E_2 \parallel \varphi T$, then $\{T, \varphi T, \xi\}$ is the Frenet frame field of γ and we have*

- 1) *if $c < 1$ then γ is biharmonic if and only if it is a geodesic;*
- 2) *if $c > 1$ then γ is proper-biharmonic if and only if it is a helix with $\kappa_1^2 = c-1$ (and $\kappa_2 = 1$).*

Remark 3.4. In dimension 3 the result was obtained by Inoguchi in [19] and explicit examples are given in [15].

Case IV ($c \neq 1$ and $g(E_2, \varphi T)$ is not constant 0, 1 or -1)

Theorem 3.5 ([16]). *Let $c \neq 1$ and γ a Legendre Frenet curve of osculating order r such that $g(E_2, \varphi T)$ is not constant 0, 1 or -1 . We have*

- 1) *if $c \leq -3$ then γ is biharmonic if and only if it is a geodesic;*
- 2) *if $c > -3$ then γ is proper-biharmonic if and only if $r \geq 4$, $\varphi T = \cos \alpha_0 E_2 + \sin \alpha_0 E_4$ and*

$$\begin{cases} \kappa_1, \kappa_2, \kappa_3 = \text{constant} > 0 \\ \kappa_1^2 + \kappa_2^2 = \frac{c+3}{4} + \frac{3(c-1)}{4} \cos^2 \alpha_0 \\ \kappa_2\kappa_3 = -\frac{3(c-1)}{8} \sin(2\alpha_0), \end{cases}$$

where $\alpha_0 \in (0, 2\pi) \setminus \{\frac{\pi}{2}, \pi, \frac{3\pi}{2}\}$ is a constant such that

$$c + 3 + 3(c-1) \cos^2 \alpha_0 > 0, \quad 3(c-1) \sin(2\alpha_0) < 0.$$

In order to obtain explicit examples of proper-biharmonic Legendre curves given by Theorem 3.1 we used the unit Euclidean sphere $\mathbb{S}^{2n+1}$ as a model of a Sasakian space form with $c = 1$ and we proved the following

Theorem 3.6 ([16]). *Let $\gamma : I \rightarrow \mathbb{S}^{2n+1}(1)$, $n \geq 2$, be a proper-biharmonic Legendre curve parametrized by arc length. Then the parametric equation of γ in the Euclidean space $\mathbb{E}^{2n+2} = (\mathbb{R}^{2n+2}, \langle \cdot, \cdot \rangle)$ is either*

$$\gamma(s) = \frac{1}{\sqrt{2}} \cos(\sqrt{2}s) e_1 + \frac{1}{\sqrt{2}} \sin(\sqrt{2}s) e_2 + \frac{1}{\sqrt{2}} e_3,$$

where $\{e_i, \mathcal{I}e_j\}$ are constant unit vectors orthogonal to each other, or

$$\gamma(s) = \frac{1}{\sqrt{2}} \cos(As) e_1 + \frac{1}{\sqrt{2}} \sin(As) e_2 + \frac{1}{\sqrt{2}} \cos(Bs) e_3 + \frac{1}{\sqrt{2}} \sin(Bs) e_4,$$

where

$$A = \sqrt{1 + \kappa_1}, \quad B = \sqrt{1 - \kappa_1}, \quad \kappa_1 \in (0, 1),$$

$\{e_i\}$ are constant unit vectors orthogonal to each other such that

$$\langle e_1, \mathcal{I}e_3 \rangle = \langle e_1, \mathcal{I}e_4 \rangle = \langle e_2, \mathcal{I}e_3 \rangle = \langle e_2, \mathcal{I}e_4 \rangle = 0$$

$$A \langle e_1, \mathcal{I}e_2 \rangle + B \langle e_3, \mathcal{I}e_4 \rangle = 0$$

and $\mathcal{I}$ is the usual complex structure on $\mathbb{R}^{2n+2}$.

Remark 3.7. For the Cases II and III we also obtained the explicit equations of proper-biharmonic Legendre curves in odd dimensional spheres endowed with the deformed Sasakian structure introduced in [27].

In [21] are introduced the complex torsions for a Frenet curve in a complex manifold. In the same way, for $\gamma : I \rightarrow N$ a Legendre Frenet curve of osculating order r in a Sasakian manifold $(N^{2n+1}, \varphi, \xi, \eta, g)$, we define the φ -torsions $\tau_{ij} = g(E_i, \varphi E_j) = -g(\varphi E_i, E_j)$, $i, j = 1, \dots, r$, $i < j$.

It is easy to see that

Proposition 3.8. *Let $\gamma : I \rightarrow N(c)$ be a proper-biharmonic Legendre Frenet curve in a Sasakian space form $N(c)$, $c \neq 1$. Then $c > -3$ and τ_{12} is constant.*

Moreover

Proposition 3.9. *If γ is a proper-biharmonic Legendre Frenet curve in a Sasakian space form $N(c)$, $c > -3$, $c \neq 1$, of osculating order $r < 4$, then it is a circle or a helix with constant φ -torsions.*

Proof. From Theorems 3.2, 3.3 and 3.5 we see that if γ is a proper-biharmonic Legendre Frenet curve of osculating order $r < 4$, then $\tau_{12} = 0$ or $\tau_{12} = \pm 1$ and, obviously, we only have to prove that when γ is a helix then τ_{13} and τ_{23} are constants.

Indeed, by using the Frenet equations of γ , we have

$$\begin{aligned} \tau_{13} &= g(E_1, \varphi E_3) = -\frac{1}{\kappa_2} g(\varphi E_1, \nabla_{E_1} E_2 + \kappa_1 E_1) = -\frac{1}{\kappa_2} g(\varphi E_1, \nabla_{E_1} E_2) \\ &= \frac{1}{\kappa_2} g(E_2, \nabla_{E_1} \varphi E_1) = \frac{1}{\kappa_2} g(E_2, \varphi \nabla_{E_1} E_1 + \xi) = 0 \end{aligned}$$

since

$$g(E_2, \xi) = \frac{1}{\kappa_1} g(\nabla_{E_1} E_1, \xi) = -\frac{1}{\kappa_1} g(E_1, \nabla_{E_1} \xi) = \frac{1}{\kappa_1} g(E_1, \varphi E_1) = 0.$$

On the other hand, it is easy to see that for any Frenet curve of osculating order 3 we have $\tau_{23} = \frac{1}{\kappa_1}(\tau'_{13} + \kappa_2 \tau_{12} + \eta(E_3))$ and

$$\begin{aligned} \eta(E_3) &= g(E_3, \xi) = \frac{1}{\kappa_2} \left(g(\nabla_{E_1} E_2, \xi) + \kappa_1 g(E_1, \xi) \right) = -\frac{1}{\kappa_2} g(E_2, \nabla_{E_1} \xi) \\ &= -\frac{1}{\kappa_2} \tau_{12}. \end{aligned}$$

In conclusion, $\tau_{23} = \frac{1}{\kappa_1}(\tau'_{13} + \kappa_2 \tau_{12} - \frac{1}{\kappa_2} \tau_{12}) = \text{constant}$.

□

Proposition 3.10. *If γ is a proper-biharmonic Legendre Frenet curve in a Sasakian space form $N(c)$ of osculating order $r = 4$, then $c \in (\frac{7}{3}, 5)$ and the curvatures of γ are*

$$\kappa_1 = \frac{\sqrt{c+3}}{2}, \quad \kappa_2 = \frac{1}{2} \sqrt{\frac{6(c-1)(5-c)}{c+3}}, \quad \kappa_3 = \frac{1}{2} \sqrt{\frac{3(c-1)(3c-7)}{c+3}}.$$

Moreover, the φ -torsions of γ are given by

$$\begin{cases} \tau_{12} = \mp \sqrt{\frac{2(5-c)}{c+3}}, & \tau_{13} = 0, & \tau_{14} = \pm \sqrt{\frac{3c-7}{c+3}}, \\ \tau_{23} = \mp \frac{3c-7}{\sqrt{3(c-1)(c+3)}}, & \tau_{24} = 0, & \tau_{34} = \pm \sqrt{\frac{2(5-c)(3c-7)}{3(c-1)(c+3)}}. \end{cases}$$

Proof. Let γ be a proper-biharmonic Legendre Frenet curve in $N(c)$ of osculating order $r = 4$. Then $c \neq 1$ and τ_{12} is different from 0, 1 or -1 . From Theorem 3.5 we have $\varphi E_1 = \cos \alpha_0 E_2 + \sin \alpha_0 E_4$. It results that

$$\tau_{12} = -\cos \alpha_0, \quad \tau_{13} = 0, \quad \tau_{14} = -\sin \alpha_0, \quad \text{and} \quad \tau_{24} = 0.$$

In order to prove that τ_{23} is constant we differentiate the expression of φE_1 along γ and using the Frenet equations we obtain

$$\begin{aligned} \nabla_{E_1} \varphi E_1 &= \cos \alpha_0 \nabla_{E_1} E_2 + \sin \alpha_0 \nabla_{E_1} E_4 \\ &= -\kappa_1 \cos \alpha_0 E_1 + (\kappa_2 \cos \alpha_0 - \kappa_3 \sin \alpha_0) E_3. \end{aligned}$$

On the other hand, $\nabla_{E_1} \varphi E_1 = \kappa_1 \varphi E_2 + \xi$ and therefore we have

$$(3.1) \quad \kappa_1 \varphi E_2 + \xi = -\kappa_1 \cos \alpha_0 E_1 + (\kappa_2 \cos \alpha_0 - \kappa_3 \sin \alpha_0) E_3.$$

We take the scalar product in (3.1) with ξ and obtain

$$(3.2) \quad (\kappa_2 \cos \alpha_0 - \kappa_3 \sin \alpha_0) \eta(E_3) = 1.$$

In the same way as in the proof of Proposition 3.9 we get

$$\begin{aligned} \eta(E_3) &= g(E_3, \xi) = \frac{1}{\kappa_2} \left(g(\nabla_{E_1} E_2, \xi) + \kappa_1 g(E_1, \xi) \right) = -\frac{1}{\kappa_2} g(E_2, \nabla_{E_1} \xi) \\ &= -\frac{1}{\kappa_2} \tau_{12} = \frac{\cos \alpha_0}{\kappa_2} \end{aligned}$$

and then, from (3.2),

$$\kappa_2 \sin \alpha_0 = -\kappa_3 \cos \alpha_0.$$

Therefore $\alpha_0 \in (\frac{\pi}{2}, \pi) \cup (\frac{3\pi}{2}, 2\pi)$.

Next, from Theorem 3.5, we have

$$\kappa_1^2 = \frac{c+3}{4}, \quad \kappa_2^2 = \frac{3(c-1)}{4} \cos^2 \alpha_0, \quad \kappa_3^2 = \frac{3(c-1)}{4} \sin^2 \alpha_0,$$

and so c must be greater than 1.

Now, we take the scalar product in (3.1) with E_3 , φE_2 and φE_4 , respectively, and we get

$$(3.3) \quad \kappa_1 \tau_{23} = -(\kappa_2 \cos \alpha_0 - \kappa_3 \sin \alpha_0) + \eta(E_3) = -\frac{\kappa_2}{\cos \alpha_0} + \frac{\cos \alpha_0}{\kappa_2}$$

$$(3.4) \quad \kappa_1 \sin^2 \alpha_0 = -(\kappa_2 \cos \alpha_0 - \kappa_3 \sin \alpha_0) \tau_{23} = -\frac{\kappa_2}{\cos \alpha_0} \tau_{23}$$

$$(3.5) \quad 0 = \kappa_1 \cos \alpha_0 \sin \alpha_0 + (\kappa_2 \cos \alpha_0 - \kappa_3 \sin \alpha_0) \tau_{34} = \kappa_1 \cos \alpha_0 \sin \alpha_0 + \frac{\kappa_2}{\cos \alpha_0} \tau_{34}.$$

and then, equations (3.3) and (3.4) lead to

$$\kappa_1^2 \sin^2 \alpha_0 = \frac{\kappa_2^2}{\cos^2 \alpha_0} - 1.$$

We come to the conclusion $\sin^2 \alpha_0 = \frac{3c-7}{c+3}$, so $c \in (\frac{7}{3}, 5)$, and then we obtain the expressions of the curvatures and the φ -torsions. $\square$

Remark 3.11. The proper-biharmonic Legendre curves given by Theorem 3.6 (for the case $c = 1$) have also constant φ -torsions.

4. A METHOD TO OBTAIN BIHARMONIC SUBMANIFOLDS IN A SASAKIAN SPACE FORM

In [16] we gave a method to obtain proper-biharmonic anti-invariant submanifolds in a Sasakian space form from proper-biharmonic integral submanifolds.

Theorem 4.1 ([16]). *Let $(N^{2n+1}, \varphi, \xi, \eta, g)$ be a strictly regular Sasakian space form with constant φ -sectional curvature c and let $\mathbf{i}: M \rightarrow N$ be an r -dimensional integral submanifold of N , $1 \leq r \leq n$. Consider*

$$F: \widetilde{M} = I \times M \rightarrow N, \quad F(t, p) = \phi_t(p) = \phi_p(t)$$

where $I = \mathbb{S}^1$ or $I = \mathbb{R}$ and $\{\phi_t\}_{t \in I}$ is the flow of the vector field ξ . Then $F: (\widetilde{M}, \widetilde{g} = dt^2 + \mathbf{i}^*g) \rightarrow N$ is a Riemannian immersion and it is proper-biharmonic if and only if M is a proper-biharmonic submanifold of N .

The previous Theorem provides a classification result for proper-biharmonic surfaces in a Sasakian space form, which are invariant under the flow-action of ξ .

Theorem 4.2 ([16]). *Let M^2 be a surface of $N^{2n+1}(c)$ invariant under the flow-action of the characteristic vector field ξ . Then M is proper-biharmonic if and only if, locally, it is given by $x(t, s) = \phi_t(\gamma(s))$, where γ is a proper-biharmonic Legendre curve.*

Also, using the standard Sasakian 3-structure on $\mathbb{S}^7$, by iteration, Theorem 4.1 leads to examples of 3-dimensional proper-biharmonic submanifolds of $\mathbb{S}^7$.

5. BIHARMONIC HOPF CYLINDERS IN A SASAKIAN SPACE FORM

Let $(N^{2n+1}, \varphi, \xi, \eta, g)$ be a strictly regular Sasakian manifold and $\bar{\mathbf{i}} : \bar{M} \rightarrow \bar{N}$ a submanifold of $\bar{N}$. Then $M = \pi^{-1}(\bar{M})$ is the Hopf cylinder over $\bar{M}$, where $\pi : N \rightarrow \bar{N} = N/\xi$ is the Boothby-Wang fibration.

In [19] the biharmonic Hopf cylinders in a 3-dimensional Sasakian space form are classified.

Theorem 5.1 ([19]). *Let $S_{\bar{\gamma}}$ be a Hopf cylinder, where $\bar{\gamma}$ is a curve in the orbit space of $N^3(c)$, parametrized by arc length. We have*

- 1) *if $c \leq 1$, then $S_{\bar{\gamma}}$ is biharmonic if and only if it is minimal;*
- 2) *if $c > 1$, then $S_{\bar{\gamma}}$ is proper-biharmonic if and only if the curvature $\bar{\kappa}$ of $\bar{\gamma}$ is constant $\bar{\kappa}^2 = c - 1$.*

In [17] we obtained a geometric characterization of biharmonic Hopf cylinders of any codimension in an arbitrary Sasakian space form. A special case of our result is the case when $\bar{M}$ is a hypersurface.

Proposition 5.2 ([17]). *If $\bar{M}$ is a hypersurface of $\bar{N}$, then $M = \pi^{-1}(\bar{M})$ is biharmonic if and only if*

$$\begin{cases} \Delta^\perp H = \left(-\|B\|^2 + \frac{c(n+1)+3n-1}{2} \right) H \\ 2 \operatorname{trace} A_{\nabla^\perp H}(\cdot) + n \operatorname{grad}(\|H\|^2) = 0, \end{cases}$$

where B , A and H are the second fundamental form of M in N , the shape operator and the mean curvature vector field, respectively, and $\nabla^\perp$ and $\Delta^\perp$ are the normal connection and Laplacian on the normal bundle of M in N .

Proposition 5.3 ([17]). *If $\bar{M}$ is a hypersurface and $\|\bar{H}\| = \text{constant} \neq 0$, then $M = \pi^{-1}(\bar{M})$ is proper-biharmonic if and only if*

$$\|\bar{B}\|^2 = \frac{c(n+1) + 3n - 5}{2}.$$

Remark 5.4. From the last result we see that there exist no proper-biharmonic hypersurfaces of constant mean curvature $M = \pi^{-1}(\bar{M})$ in $N(c)$ if $c \leq \frac{5-3n}{n+1}$, which implies that such hypersurfaces do not exist if $c \leq -3$, whatever the dimension of N is.

From now on we shall consider $c > -3$.

In [26] Takagi classified all homogeneous real hypersurfaces in the complex projective space $\mathbb{C}P^n$, $n > 1$, and found five types of such hypersurfaces (see also [23]). The first type (with subtypes A1 and A2) are described in the following.

We shall consider $u \in (0, \frac{\pi}{2})$ and r a positive constant given by $\frac{1}{r^2} = \frac{c+3}{4}$.

Theorem 5.5 ([26]). *The geodesic spheres (Type A1) in complex projective space $\mathbb{C}P^n(c+3)$ have two distinct principal curvatures: $\lambda_2 = \frac{1}{r} \cot u$ of multiplicity $2n-2$ and $a = \frac{2}{r} \cot(2u)$ of multiplicity 1.*

Theorem 5.6 ([26]). *The hypersurfaces of Type A2 in complex projective space $\mathbb{C}P^n(c+3)$ have three distinct principal curvatures: $\lambda_1 = -\frac{1}{r} \tan u$ of multiplicity $2p$, $\lambda_2 = \frac{1}{r} \cot u$ of multiplicity $2q$, and $a = \frac{2}{r} \cot(2u)$ of multiplicity 1, where $p > 0$, $q > 0$, and $p + q = n - 1$.*

We note that if $c = 1$ and $\bar{M}$ is of type A1 or A2 then $\pi^{-1}(\bar{M}) = \mathbb{S}^1(\cos u) \times \mathbb{S}^{2n-1}(\sin u) \subset \mathbb{S}^{2n+1}$ or $\pi^{-1}(\bar{M}) = \mathbb{S}^{2p+1}(\cos u) \times \mathbb{S}^{2q+1}(\sin u)$, respectively.

By using Takagi's result we classified in [17] the biharmonic Hopf cylinders $M = \pi^{-1}(\bar{M})$ in a Sasakian space form N^{2n+1} over homogeneous real hypersurfaces in $\mathbb{C}P^n$, $n > 1$.

Theorem 5.7 ([17]). *Let $M = \pi^{-1}(\bar{M})$ be the Hopf cylinder over $\bar{M}$.*

1) *If $\bar{M}$ is of Type A1, then M is proper-biharmonic if and only if either*

- a) $c = 1$ and $\tan^2 u = 1$, or
- b) $c \in \left[\frac{-3n^2+2n+1+8\sqrt{2n-1}}{n^2+2n+5}, +\infty \right) \setminus \{1\}$ and

$$\tan^2 u = n + \frac{2c-2}{c+3} \pm \frac{\sqrt{c^2(n^2+2n+5)+2c(3n^2-2n-1)+9n^2-30n+13}}{c+3}.$$

2) *If $\bar{M}$ is of Type A2, then M is proper-biharmonic if and only if either*

- a) $c = 1$, $\tan^2 u = 1$ and $p \neq q$, or
- b) $c \in \left[\frac{-3(p-q)^2-4n+4+8\sqrt{(2p+1)(2q+1)}}{(p-q)^2+4n+4}, +\infty \right) \setminus \{1\}$ and

$$\tan^2 u = \frac{n}{2p+1} + \frac{2c-2}{(c+3)(2p+1)} \pm \frac{\sqrt{c^2((p-q)^2+4n+4)+2c(3(p-q)^2+4n-4)+9(p-q)^2-12n+4}}{(c+3)(2p+1)}.$$

Theorem 5.8 ([17]). *There are no proper-biharmonic hypersurfaces $M = \pi^{-1}(\bar{M})$ when $\bar{M}$ is a hypersurface of Type B, C, D or E in the complex projective space $\mathbb{C}P^n(c+3)$.*

ACKNOWLEDGEMENTS

The authors were partially supported by the Grant CEEEX, ET, 5871/2006 and by the Grant CEEEX, ET, 5883/2006, Romania.

The first author would like to thank to the organizers, especially to Professor I. Mladenov, for the Conference Grant.

REFERENCES

- [1] K. Arslan, R. Ezentas, C. Murathan and T. Sasahara, *Biharmonic Anti-invariant Submanifolds in Sasakian Space Forms*, Beitrge Algebra Geom., **48** (1)(2007), 191–207.
- [2] A. Arvanitoyeorgos, F. Defever, G. Kaimakamis and V.J. Papantoniou, *Biharmonic Lorentz Hypersurfaces in E_1^4* , Pacific J. Math., **229** (2007), 293–305.
- [3] A. Arvanitoyeorgos, F. Defever and G. Kaimakamis, *Hypersurfaces of E_s^4 with Proper Mean Curvature Vector*, J. Math. Soc. Japan, **59** (2007), 797–809.
- [4] A. Balmuş, S. Montaldo and C. Oniciuc, *Classification Results for Biharmonic Submanifolds in Spheres*, Israel J. Math., to appear.
- [5] A. Balmuş, S. Montaldo and C. Oniciuc, *Biharmonic Hypersurfaces in 4-Dimensional Space Forms*, Math. Nachr., to appear.
- [6] D.E. Blair, *Riemannian Geometry of Contact and Symplectic Manifolds*, Birkhäuser Boston, Progress in Mathematics, Volume 203, 2002.
- [7] R. Caddeo, S. Montaldo and C. Oniciuc, *Biharmonic Submanifolds of $\mathbb{S}^3$* , Internat. J. Math., **12** (2001), 867–876.
- [8] R. Caddeo, S. Montaldo and C. Oniciuc, *Biharmonic Submanifolds in Spheres*, Israel J. Math., **130** (2002), 109–123.

- [9] R. Caddeo, S. Montaldo and P. Piu, *Biharmonic Curves on a Surface*, Rend. Mat. Appl., (7) **21** (2001), 143–157.
- [10] B.Y. Chen, *A Report on Submanifolds of Finite Type*, Soochow J. Math., **22** (1996), 117–337.
- [11] B.Y. Chen, *Classification of Marginally Trapped Lorentzian Flat Surfaces in $\mathbb{E}_2^4$ and Its Application to Biharmonic Surfaces*, J. Math. Anal. Appl., **340** (2008), 861–875.
- [12] J.T. Cho, J. Inoguchi and J.-E. Lee, *Biharmonic Curves in 3-Dimensional Sasakian Space Form*, Ann. Math. Pura Appl., **186** (2007), 685–701.
- [13] I. Dimitric, *Submanifolds of $\mathbb{E}^m$ with Harmonic Mean Curvature Vector*, Bull. Inst. Math. Acad. Sinica, **20** (1992), 53–65.
- [14] J. Eells and J.H. Sampson, *Harmonic Mappings of Riemannian Manifolds*, Amer. J. Math., **86** (1964), 109–160.
- [15] D. Fetcu and C. Oniciuc, *Explicit Formulas for Biharmonic Submanifolds in Non-Euclidean 3-Spheres*, Abh. Math. Sem. Univ. Hamburg, **77** (2007), 179–190.
- [16] D. Fetcu and C. Oniciuc, *Explicit Formulas for Biharmonic Submanifolds in Sasakian Space Forms*, [arXiv:math.DG/0706.4160v1](https://arxiv.org/abs/math/0706.4160v1).
- [17] D. Fetcu and C. Oniciuc, *Biharmonic Hypersurfaces in Sasakian Space Forms*, Preprint.
- [18] T. Ichiyama, J. Inoguchi and H. Urakawa, *Bi-harmonic Maps and Bi-Yang-Mills Fields*, Note Mat., to appear.
- [19] J. Inoguchi, *Submanifolds with Harmonic Mean Curvature in Contact 3-manifolds*, Colloq. Math., **100** (2004), 163–179.
- [20] G.Y. Jiang, *2-harmonic Maps and Their First and Second Variational Formulas*, Chinese Ann. Math. Ser. A, **7** (4)(1986), 389–402.
- [21] S. Maeda and Y. Ohnita, *Helical Geodesic Immersions into Complex Space Forms*, Geom. Dedicata, **30** (1989), 93–114.
- [22] S. Montaldo and C. Oniciuc, *A Short Survey on Biharmonic Maps Between Riemannian Manifolds*, Rev. Un. Mat. Argentina, **47** (2)(2006), 1–22.
- [23] R. Niebergall and P.J. Ryan, *Real Hypersurfaces in Complex Space Forms*, Tight and Taut Submanifolds, MSRI Publications, **32** (1997), 233–305.
- [24] K. Ogiue, *On Fiberings of Almost Contact Manifolds*, Kōdai Math. Sem. Rep., **17** (1965), 53–62.
- [25] T. Sasahara, *Legendre Surfaces in Sasakian Space Forms Whose Mean Curvature Vectors Are Eigenvectors*, Publ. Math. Debrecen, **67** (3–4)(2005), 285–303.
- [26] R. Takagi, *On Homogeneous Real Hypersurfaces in a Complex Projective Space*, Osaka J. Math., **10** (1973), 495–506.
- [27] S. Tanno, *The Topology of Contact Riemannian Manifolds*, Ill. J. Math., **12** (1968), 700–717.
- [28] The Bibliography of Biharmonic Maps. <http://beltrami.sc.unica.it/biharmonic/>.
- [29] W. Zhang, *New Examples of Biharmonic Submanifolds in $\mathbb{C}P^n$ and $\mathbb{S}^{2n+1}$* , [arXiv:math.DG/0705396v1](https://arxiv.org/abs/math/0705396v1).

DEPARTMENT OF MATHEMATICS, "GH. ASACHI" TECHNICAL UNIVERSITY OF IASI, BD. CAROL I NO. 11, 700506 IASI, ROMANIA

E-mail address: dfetcu@math.tuiasi.ro

FACULTY OF MATHEMATICS, "AL.I. CUZA" UNIVERSITY OF IASI, BD. CAROL I NO. 11, 700506 IASI, ROMANIA

E-mail address: oniciucc@uaic.ro